



## Potential use of mud clam (*Geloina coaxans*) in producing sauce with papaya crude extraction as a protein hydrolysing agent

Hitihami Mudiyanselage Sarath Mahinda Wijerathna<sup>1</sup> • Kumudu Radampola<sup>2</sup> • Hewa Waduge Cyril<sup>3</sup>

<sup>1</sup> Department of Aquaculture and Aquatic Resources Management, University College of Anuradhapura, University of Vocational Technology, Sri Lanka

<sup>2</sup> Department of Fisheries and Aquaculture, Faculty of Fisheries and Marine sciences & Technology, University of Ruhuna, Matara, Sri Lanka

<sup>3</sup> Department of Animal science, Faculty of Agriculture, University of Peradeniya, Peradeniya, Sri Lanka

### Correspondence

Hitihami Mudiyanselage Sarath Mahinda Wijerathna; Department of Aquaculture and Aquatic Resources Management, University College of Anuradhapura, University of Vocational Technology, Sri Lanka

[sarawijem@gmail.com](mailto:sarawijem@gmail.com)

### Manuscript history

Received 3 August 2020 | Revised 10 March 2021 | Accepted 12 March 2021 | Published online 17 March 2021

### Citation

Wijerathna HMSM, Radampola K, Cyril HW (2021) Potential use of mud clam (*Geloina coaxans*) in producing sauce with papaya crude extraction as a protein hydrolysing agent. Journal of Fisheries 9(1): 91402. DOI: 10.17017/j.fish.269

### Abstract

Mud clam (*Geloina coaxans*) are underutilised food source due to the lack of consumer preference in Sri Lanka. Hence, this study was conducted to produce clam sauce as a value added product using the muscle of mud clams by means of accelerated fermentation method. Specimens were collected from Tambalagam Bay, Sri Lanka. Shell length, height, inflation, total weight with shell and without shell were  $6.3 \pm 0.4$  cm,  $5.3 \pm 0.3$  cm,  $3.5 \pm 0.3$  cm,  $83.1 \pm 13.4$  g and  $14.9 \pm 1.3$  g respectively. The extracted mean meat yield was  $14.9 \pm 1.3\%$  per mud clam. Moisture, crude protein, crude lipid and ash (dry weight) content of raw meat were  $80.45 \pm 0.89\%$ ,  $64.14 \pm 0.96\%$ ,  $3.55 \pm 0.39\%$  and  $7.54 \pm 0.61\%$  respectively. Final sauce product shows liquid yield, energy value,  $^0$ Brix value, pH, % NaCl, total nitrogen, moisture and ash content as  $98.3 \pm 5.5$  ml  $100\text{g}^{-1}$ ,  $2124 \pm 133$  J  $\text{g}^{-1}$ ,  $24.3 \pm 0.9\%$ ,  $5.02 \pm 0.04$ ,  $14.53 \pm 0.27\%$ ,  $0.27 \pm 0.01\%$ ,  $74.06 \pm 0.56\%$  and  $19.66 \pm 1.99\%$  respectively. The study concluded that the mud clam meat is a possible candidate as a raw material for the production of clam sauce.

**Keywords:** Accelerated fermentation; clam sauce; *Geloina coaxans*

### 1 | INTRODUCTION

Bivalve exploitation and cultivation for food, ornamentation and pearls are worldwide practices for decades and farming for food is a multi-billion-dollar industry worldwide (Wijsman *et al.* 2019). Despite the abundance of edible bivalve species in most of the coastal zones of Sri Lanka (Chinivasagam 1989) value addition of bivalves is not much popular (Silva *et al.* 2013). Edible bivalves such as *Marcia hiantina*, *M. opima*, *Gastrarium tumidum* and *Meretrix casta* are commonly found in Puttalam Lagoon and Dutch Canal in the west coast of Sri Lanka (Kithsiri *et*

*al.* 2004; Jayawikrema and Wijeyaratne 2009). Furthermore, diverse edible clams, mussels and oyster species are also abundant in the Tambalagam Bay in Trincomalee district along the east coast of Sri Lanka (Nafees *et al.* 2009). The mud clam *Geloina coaxans* is the most common bivalve species in Tambalagam Bay area.

Bivalves are a good source of protein, vitamins (e.g. vitamin C and Vitamin B<sub>12</sub>) and minerals (e.g. iron, manganese, phosphorus, selenium and zinc) and fatty acids for humans (Shanmugam *et al.* 2007; Patrick 2012). Despite the fact that it is a luxury food item in many foreign

countries, it is not much popular among Sri Lankans (Silva et al. 2013). Hence, local consumption could be increased by encouraging people through promoting and marketing it as a value added product. Sauce is one of the value added products, which is produced by using different kind of raw materials such as fish (marine and freshwater) and oysters, chilly, tomato and soybean. In many South East Asian countries, fish is mixed with salt and keep for one to two years for fermentation to produce their traditional fish sauce (Hariono et al. 2005). However, enzymes can be used as a protein hydrolysing agent to speed up the fermentation process. Papain is a very effective protein hydrolysing enzyme that can be extracted from papaya (*Carica papaya*) latex, unripe fruit, leaves and roots (Hitesh et al. 2012). The present study was carried out to investigate the possibility of using papaya crude extract as a source of papain enzyme to hydrolyse the protein in mud clam muscle to produce clam sauce through accelerated fermentation method.

## 2 | METHODOLOGY

One hundred mud clams (*G. coaxans*) were purchased from a fisherman in Tambalagam Bay, Kinniya, Sri Lanka. Biometric parameters (shell height, length, inflation and weight of the whole animal with the shell) were measured for each individual. Afterwards, shell of each bivalve was removed and weight of the muscle was recorded.

Separated muscle parts of the clam individuals were immediately sealed in polythene bags and kept in a polystyrene box with ice to retain freshness during the transportation to the laboratory of the Department of Animal Science, University of Peradeniya. Soon after the arrival, proximate composition of raw muscle was analysed by means of standard methods (AOAC 1995). Bivalve sauce was prepared according to the method described by Chuapoeuk and Raksakulathai (1992), with some modifications in which unripe papaya crude extraction was used as a Protein hydrolysing agent. Hush and seeds of unripe papaya fruit were removed and the fruit was cut into small pieces, which were ground into a pulp. The papaya pulp was filtered through a linen cloth to separate papaya crude extract and kept it inside a refrigerator in a lid closed glass bottle. Cleaned, frozen clam muscles were sliced and ground to make a pulp. For the fermentation process 100 g of mud clam muscle pulp, 40 g of table salt (NaCl) and 60 g of papaya crude extract were mixed thoroughly in a glass bottle. Then, the bottle was closed with a para film and sealed properly and kept it at 60°C in an oven for 10 days until fermentation. After 10 days, the mixture was filtered through a Whatman No. 4 filter paper into a glass bottle and a dark brownish colour filtrate was obtained as the clam sauce. The volume of the filtrate was measured and liquid yield of the clam sauce was determined. Then, the prepared sauce was subjected to analyse for proximate composition, pH, salt content

(Volhard method), °Brix value and energy value (Chuapoeuk and Raksakulathai 1992; AOAC 1995; Hariono et al. 2005; Muangthai et al. 2009).

## 3 | RESULT AND DISCUSSION

The biometric measurements and percentage of total yield of mud clam which were harvested from Tambalagam Bay, Kinniya, Sri Lanka are presented in Table 1. Percentage mean total yield of the mud clam was  $18.1 \pm 1.8\%$ . *Geloina coaxans* has a thick and heavy shell and thus, the weight of flesh obtained from mud clam is very low when compared to its total weight.

**TABLE 1** Biometric parameters of *Geloina coaxans* ( $n = 100$ ).

Biometric parameters	Mean ( $\pm SD$ )	Range
Shell length (cm)	$6.3 \pm 0.4$	5.6 – 7.2
Shell height (cm)	$5.3 \pm 0.3$	4.7 – 6.1
Shell inflation (cm)	$3.5 \pm 0.3$	3.0 – 4.3
Total weight (with shell) (g)	$83.1 \pm 13.4$	50.0 – 120.2
Weight (without shell) (g)	$14.9 \pm 1.3$	12.4 – 17.5
Total yield (%)	$18.1 \pm 1.8$	13.7 – 22.7

Proximate composition of *G. coaxans* muscle is shown in Table 2. Asha et al. (2014) reported that the crude protein and crude lipid content in muscles of *Crassostrea madrasensis*, a common bivalve species use in commercial oyster sauce industry, as  $9.41 \pm 0.85\%$  and  $3.25 \pm 0.32\%$  (wet weight basis) respectively. In the present study, *G. coaxan* contains higher amount of protein (12.62%) and lower amount of lipid (0.69%) when compared to *C. madrasensis*. Therefore, mud clam can be used as a low lipid and high protein source for human consumption.

**TABLE 2** Proximate composition of *Geloina coaxans* (mean  $\pm SD$ ,  $n = 3$ ).

Proximate composition	Dry weight basis (%)	Wet weight basis (%)
Moisture	–	$80.45 \pm 0.89$
Crude protein	$64.14 \pm 0.96$	$12.62 \pm 0.19$
Crude lipid	$3.55 \pm 0.39$	$0.69 \pm 0.08$
Ash	$7.54 \pm 0.61$	$1.47 \pm 0.12$

*Geloina coaxans* is a low-priced and commonly available bivalve in the Tambalagam Bay. Therefore, it was used as the raw material to investigate the potential of production of clam sauce by means of accelerated fermentation method in this study. Papaya crude extract was used as an inexpensive locally available raw material to hydrolyse the protein in bivalve muscles during the fermentation process. Papaya crude extract contains cysteine protease, also known as papain or papaya proteinase which has the ability to hydrolyse the protein into amino acids (Hitesh et al. 2012). Accelerated fermenta-

tion process with papaya crude extract resulted in 98.3 ml 100g<sup>-1</sup> of liquid yield of clam sauce (Table 3). However, Hariono et al. (2005) have extracted only 45 ml of fish sauce from 100 g of round scad (*Decapterus macrosoma*) and hard tail (*Megalaspis cordyla*) fish meats. Hence, considerably higher amount of liquid yield was produced in this study when compare with the previous studies.

The total soluble solid content (°Brix) value depends on the amount of free amino acids and small peptides in the sauce (Hariono et al. 2005) which was 24.3% for *G. coaxans*. However, this value is 41.5% for commercial fish sauce (Hariono et al. 2005) which is higher than the °Brix value of clam sauce produced in this study. This was also the case for total-N content in the mud clam sauce in the present study (0.27%; Table 3) whereas the commercial fish sauce comprises 1.95% (Hariono et al. 2005). However, commercial sauces are usually mixed with different additives to enhance the flavour, texture and nutritional values. Especially monosodium glutamate (MSG) is used as a chemical additive to enhance flavour and it is a sodium salt of the naturally occurring non-essential amino acids (Kingsley et al. 2013). As a result, commercial products show high values of °Brix and total-N than the clam sauce produced in the present study which may be due to no use of any preservatives or additives in the present study.

**TABLE 3** Liquid yield, energy content, °Brix value, chemical and proximate composition (wet weight basis) of the sauce produced from *Geloina coaxans* (mean ± SD, n = 3).

Properties	This study	Commercial fish sauce
Liquid yield (ml 100g <sup>-1</sup> )	98.3 ± 5.5	–
Energy value (J g <sup>-1</sup> )	2124 ± 133	–
Brix (%)	24.3 ± 0.9	41.5 (Hariono et al. 2005)
pH	5.02 ± 0.04	5 – 6 (Puat et al. 2015)
NaCl (%)	14.53 ± 0.27	19 – 20 (Puat et al. 2015)
Total N (%)	0.27 ± 0.01	>2.0 (Puat et al. 2015)
Moisture (%)	74.06 ± 0.56	–
Ash (%)	19.66 ± 1.99	–

Prepared clam sauce was slightly acidic with pH of 5.02±0.09 which roughly indicates the adequacy of fermentation (Hariono et al. 2005). However, the pH value of present experiment was laid on the acceptable pH range (5 – 6) which is described in Thai industrial standard for fish sauce (Puat et al. 2015). The salt content of mud clam sauce was 14.53%; slightly similar result was obtained for oyster sauce (Chuapoeuk and Raksakulthai 1992) in which 14.6% and 13.5% of salt contents were recorded when papain and bromelain were used to produce the sauce. Salt content is important in sauce production as it can prevent the spoilage during the fermenta-

tion of product (Shiau and Chai 1999).

Present study revealed that *G. coaxans* is a possible candidate to produce clam sauce by means of accelerated fermentation method with papaya crude extract as the protein hydrolysing agent. Moreover, this method can be used as a low cost method to prepare mud clam sauce and further research is needed to develop this method which can be easily used by local fishermen to prepare sauce as a value addition product of mud clam.

## ACKNOWLEDGEMENTS

Authors are grateful to the Department of Animal Sciences, University of Peradeniya for providing laboratory facilities for the analysis.

## CONFLICT OF INTEREST

The authors declare no conflict of interest.

## AUTHORS' CONTRIBUTION

HMSM conceptualisation, data collection, data analysis and original manuscript preparation; KR supervision, manuscript review and editing; HWC supervision.

## DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available on request from the corresponding author.

## REFERENCES

- AOAC (1995) Official methods of analysis of the official Association of Official Analytical Chemists, 16th edition. Association of Official Analytical Chemists. Arlington, Virginia, USA.
- Asha K, Anandan R, Mathew S, Lakshmanan P (2014) *Biochemical profile of oyster Crassostrea madrasensis and its nutritional attributes*. The Egyptian Journal of Aquatic Research 40(1): 35–41.
- Chinivasagam HN (1989) Status of sanitation and marketing of mollusc in Sri Lanka. Regional seafarming development and demonstration project, Network of Aquaculture Centres in Asia, Bangkok, Thailand.
- Chuapoeuk P, Raksakulthai N (1992) Use of papain and bromelain in the production of oyster sauce. Asian Food Journal 7: 196–199.
- Hariono I, Yeap SE, Kok TN, Ang GT (2005) Use of Koji and protease in fish sauce fermentation. Singapore Journal of Primary Industries 32: 19–29.
- Hitesh P, Manojbhai NB, Mayuri AB, Ashvinkumar DD, Kiranben VD (2012) Extraction and application of papain enzyme on degradation of drug. International Journal of Pharmacy and Biological Sciences 2(3): 113–115.
- Jayawikrema EM, Wijeyaratne MJS (2009) *Distribution and population dynamics of the edible bivalve species Meretrix casta (Chemnitz) in the Dutch Canal of Sri Lanka*. Sri Lankan Journal of Aquatic science 14:

29–44.

- Kingsley OA, Jacks TW, Amaza DS, Peters TM, Otong ES (2013) The effect of monosodium glutamate (MSG) on the gross weight of the heart of albino rats. Scholars Journal of Applied Medical Sciences 1(2): 44–47.
- Kithsiri HMP, Wijerathne MJS, Amarasinghe US (2004) Population dynamics of Three commercially important bivalves species (Family: Veneridae) in Puttalam Lagoon and Dutch Bay, Sri Lanka. Sri Lanka Journal of Aquatic Sciences 9(1): 13–29.
- Muangthai P, Upajak P, Suwunna P, Patumpai W (2009) Development of healthy soy sauce from pigeon pea and soybean. Asian Journal of Food and Agro-Industry 2(3): 291–301.
- Nafees MSM, Athauda ARSB, Edirisinghe U (2009) Impacts of tsunami and security situations on marine fisheries of Kinniya, Trincomalee. Tropical Agricultural Research 21(1): 80–88.
- Patrick K (2012) Nutritional delights: clams, oysters, mussels and scallops. AlignLife. [www.alignlife.com/articles/food/nutritional\\_delights\\_clams\\_oysters\\_mussels\\_and\\_scallops](http://www.alignlife.com/articles/food/nutritional_delights_clams_oysters_mussels_and_scallops). Accessed on 20 July 2020.
- Puat SNA, Huda N, Abdullah WNW (2015) Chemical Composition and protein quality of fish sauces (Kecap Ikan and Nampla). Asia Pacific Journal of Sustainable Agriculture Food and Energy 3(2): 2–9.
- Shanmugam A, Palpandi C, Sambasivam S (2007) Some valuable fatty acids exposed from wedge clam *Donax cuneatus* (Linnaeus). African Journal of Biochemistry Research 1(2): 14–18.
- Shiau CY, Chai TJ (1999) Protein recovered from oyster wash water by ultra filtration and their utilization as oyster sauce through fermentation. Journal of Marine Science and Technology 7(2): 110–116.
- Silva EIL, Katupotha J, Amarasinghe O, Manthrithilake H, Ariyaratna R (2013) Lagoons of Sri Lanka: from the origins to the present. International Water Management Institute, Colombo.
- Wijsman JWM, Troost K, Fang J, Roncarati A (2019) Global production of marine bivalves. Trends and challenges. pp. 7–26. In: Smaal A, Ferreira J, Grant J, Petersen J, Strand Ø. (Eds) Goods and services of marine bivalves. Springer, Cham.



HMSM Wijerathna <https://orcid.org/0000-0002-2153-3143>  
K Radampola <https://orcid.org/0000-0003-0292-4576>