


Microstructures of barbels and bony plates of *Satyrichthys laticeps* (Actinopterygii: Peristediidae)

Arin Ngamniyom

Faculty of Environmental Culture and Eco-tourism, Srinakharinwirot University, Bangkok 10110, Thailand

Correspondence

Arin Ngamniyom; Faculty of Environmental Culture and Eco-tourism, Srinakharinwirot University, Bangkok 10110, Thailand

 arin@g.swu.ac.th

Manuscript history

Received 13 August 2019 | Revised 5 May 2020 | Accepted 6 May 2020 | Published online 9 May 2020

Citation

Ngamniyom A (2020) Microstructures of barbels and bony plates of *Satyrichthys laticeps* (Actinopterygii: Peristediidae). Journal of Fisheries 8(2): 850–853.

Abstract

Barbels are sensory-appendage near the mouth of many fishes including Cyprinidae, Ictaluridae and Peristediidae and commonly used for identification of the species. Similarly scales and bony plates of fishes also being used in taxonomic studies. *Satyrichthys laticeps* commonly called armoured searobins, is a deep-sea peristediid that distributed in the Indo-West Pacific region of the world. However, surface structures of has been unknown. The aim of this study is to reveal the microstructures of barbels and bony plates of *S. laticeps*, occurring in the Andaman Sea of Thailand, for the first time by using a scanning electron microscopy method. The chin barbels were characterised by a pebble-like surface whereas lip barbels resemble wooden texture. Taste bud-like organs were only visible in chin barbels. Bony plates from heads contained small semispheres around spine. In dorsolateral of posterior body, pits appeared at the apices, and semispherical buds were present in the middle and basal parts of tubercles. The study results may be important in understanding the physiology in peristediid fish.

Keywords: Armoured searobins; *Satyrichthys laticeps*; peristediids; SEM; microstructures

1 | INTRODUCTION

In teleosts, barbel is slender process and an integumentary sensory organ of fishes; examples include carps, catfishes and goatfishes (Grover-Johnson and Farbman 1976; Lombarte and Aguirre 1997; LeClair and Topczewski 2010). Similar to scale of fishes bony plates of varying structures are also commonly found in some fish groups such as searobins, boxfish, sturgeon and seahorse that are used to defend the body against the smashing, scratching or predator attack (Porter *et al.* 2013; Qu *et al.* 2013; Yang *et al.* 2013, 2015). Porter *et al.* (2013) described that bio-inspired design from nature has led to a

novel study of fracture-resistant structures.

Armoured searobins are peristediid teleosts belonging to the order Scorpaeniformes, an order comprising mostly marine deep-sea fish (Kawai *et al.* 2008; Van Oijen *et al.* 2013; Kawai 2016). *Satyrichthys laticeps* (Schlegel, 1852), a species of the family Peristediidae, is widely distributed in the tropical regions of the Indo-West Pacific including the Andaman Sea of Thailand (Kawai 2013, 2014; Pogoreutz *et al.* 2014). Recently morphological descriptions of *S. laticeps* from the deep-sea habitats of the Andaman Sea in the eastern Indian Ocean have been described by Kawai *et al.* (2017). Although the barbels are dominant

character in peristediids but its microstructures is still unknown. Therefore the aims of this work are to study the microstructures of barbels and some bony plates of *S. laticeps* from the Andaman Sea of Thailand.

2 | METHODOLOGY

In September 2017, 14 specimens of *S. laticeps*, including six male and eight females, were captured from the Andaman Sea of Thailand with the help of professional fishermen using a bottom trawl net. The fish individuals were euthanized by administering an overdose of a MS-222. Specimens were identified using the key for the peristediid genus *Satyrichthys* (after Kawai 2013). Barbels were cut from the lips and chins, and pieces of bony plates were stripped from the fish bodies (Figure 1). In scanning electron microscope (SEM), tissues and pieces were immediately fixed in 4% glutaraldehyde with 0.1 M phosphate buffer (pH 7.4) and left overnight at 25°C. Specimens were washed with 0.1 M phosphate buffer and post-fixed in 1% OsO₄ in the same buffer at 25°C for 60 minutes. Following OsO₄ fixation samples were rinsed with distilled water and dehydrated using an ethanol dehydration series (50, 60, 70, 80, 90, 95 and 100%) for 10 minutes in each concentration. Specimens were dried using a critical point drying apparatus surrounded by high pressure of CO₂, mounted on metal stubs with carbon tape, coated with gold using a sputter coater, analysed by a HITACHI

SU-8010 SEM, and imaged at 10 kV.

3 | RESULTS AND DISCUSSION

The mean (\pm SD) total length including the stick-rostral projection of male armoured searobins was 30.1 ± 8.6 cm (range: 24 – 47.4 cm; $n = 6$) whereas for female the mean length was 35.7 ± 9.2 cm (25.0 – 47.0 cm; $n = 8$). The SEM analysis of barbels, at high-magnification, showed that the surface of chin barbels are crumpled and covered by a pebble-like surface texture (Figures 1B – 1C). Some areas lip barbel surfaces were rough and had a wooden pattern and rest areas had a fold-like network alternating with wrinkles (Figures 1D – 1G). The surface of the branched end of the lip-barbel was covered by a narrow network of folds and wrinkles (Figures 1H – 1I). The rostral projection had margin on surfaces that are smooth and contained short digitiform tubercles with small branched prongs at the tips (Figure 1J). This study showed that the structure of chin barbels differed from lip barbels in many respects. Hussain and Bordoloi (2018) reported that the surface morphology of barbels in *Garra* spp. was peculiar and may aid the fish in adapting to their habitats. Kiyohara *et al.* (2002) described how the goatfish (*Parupeneus* spp.) uses the long chin barbels to scan for prey along the sea bottom. Therefore, sensory chin barbels of *S. laticeps* may play key role in searching food or catching prey in deep sea habitats.




FIGURE 1 Scanning electron microscope (SEM) photomicrography of barbels (B – I) and bony plates (J – U) of *Satyrichthys laticeps*. Overall locations are presented in A. B, chin barbel; C, high-magnification view of B; D, lip barbel; E, high-magnification view of D; F, branched parts of lip barbel; G, high-magnification view of F; H, branched lip barbel at posterior end; I, high-magnification view of H; J, bony plate of snout; K, plate of lateral head; L, bony plate of head; M, high magnification view of L; N, dorsal spine; O, high magnification view of short spine of N; P, plate of dorsal body; Q, high-magnification view of P; R, plate of lateral body; S, high-magnification view of R; and U, overlapping surfaces of the posterior region of bony plate.

The SEM analysis of bony plates showed that the stout symmetric nodes and smooth surfaces at the tip and distal areas of the lateral side of head (Figure 1K). Surfaces of the frontal and parietal plates were decorated with cone-shaped tubercles. Some buds were found around the base of those tubercles (Figures 1L – 1M). In dorsal view, near the spines, cone-shaped tubercles were positioned on the surfaces of bony plates. Tubercles with small buds were quite spherical and smooth (Figure 1N – 1O). In the dorsolateral areas of the posterior part of the body, there were cone-shaped tubercles that appeared to be pitted at the apices and spherical buds were present in the middle and basal parts of the tubercles. Cone-shaped tubercles were also observed on the bony-plate surfaces along the posterior part of the body (Figures 1P – 1Q). For the bony plates below the antrorse spines, the cone-shaped tubercles were stout and arranged in alternating rows with hollows. Tubercles were broadly covered by small pits. Along the surfaces of overlapping plates there were hollow areas without any tubercles (Figures 1R – 1U). Members of the family Peristediidae commonly live in the deep sea and are characterised by hard bony plates covering the body and rostral projections (Nelson 2006). Surface structure of bony plates of *S. laticeps* resembles that they may be related to their environment and may help in the movement, foraging and maintaining balance in the depth sea.

This study depicts the ultrastructure of barbels and bony plates of *S. laticeps*, a deep-sea peristediid, from the Andaman Sea of Thailand. The results are believed to have contributed to the knowledge about bony structure in Peristediidae that may be of help in understanding the biology of armoured searobins.

ACKNOWLEDGEMENTS

The author deeply thanks Miss Ornjira Tachaworarak, Department of Microbiology, Faculty of Sciences, Srinakharinwirot University, for her help in searching relevant literature.

CONFLICT OF INTEREST

The author declares no conflict of interest.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available on request from the corresponding author.

REFERENCES


- Grover-Johnson N, Farbman AI (1976) [Fine structure of taste buds in the barbel of the catfish, *Ictalurus punctatus*](#). Cell and Tissue Research 28: 395–403.
- Hussain JF, Bordoloi S (2018) [Adaptive modifications in four fish species of the genus *Garra* \(Teleostei; Cyprinidae\) in Basistha River, Assam, India](#). Microscopy and Microanalysis 24: 310–317.
- Kawai T (2013) [Revision of the peristediid genus *Satyrichthys* \(Actinopterygii: Teleostei\) with the description of a new species, *S. milleri* sp. nov.](#) Zootaxa 3635: 419–438.
- Kawai T (2014) [Satyrichthys kikingeri Pogoreutz, Vitecek & Ahnelt, 2013, a junior synonym of *Satyrichthys laticeps* \(Schlegel, 1852\) \(Actinopterygii: Teleostei: Peristediidae\)](#). Zootaxa 19(1): 135–140.
- Kawai T (2016) [Peristedion richardsi sp. nov. \(Actinopterygii: Teleostei: Peristediidae\) from Indonesian waters, with synonymy between *Peristedion riversandersoni* Alcock, 1894 and *Peristedion nierstraszi* Weber, 1913](#). Zootaxa 4171(2): 335–346.
- Kawai T, Nakaya K, Séret B (2008) [A new armored searobin *Paraheminodus longirostralis* \(Teleostei: Peristediidae\) from New Caledonia](#). Ichthyological Research 55: 374–378.
- Kawai T, Tashiro F, Imamura H, Aungtonya C (2017) Deep-sea fishes from the Andaman Sea by R/V Chakratong Tongyai during 1996–2000. Part 1: the order Scorpaeniformes. Phuket Marine Biological Center Research Bulletin 74: 23–32.
- Kiyohara S, Sakata Y, Yoshitomi T, Tsukahara J (2002) [The 'goatee' of goatfish: innervation of taste buds in the barbels and their representation in the brain](#). Proceedings of the Royal Society B: Biological Sciences 269: 1773–1780.
- LeClair EE, Topczewski J (2010) [Development and regeneration of the zebrafish maxillary barbel: a novel study system for vertebrate tissue growth and repair](#). PLoS One 15: e8737.
- Lombarte A, Aguirre H (1997) Quantitative differences in the chemoreceptor systems in the barbels of two species of Mullidae (*Mullus surmuletus* and *M. barbatus*) with different bottom habitats. Marine Ecology Progress Series 150: 57–64.
- Nelson JS (2006) Fishes of the world, fourth edition. John Wiley and Sons, New York.
- Pogoreutz C, Vitecek S, Ahnelt H (2014) [First record of *Satyrichthys laticeps* and second record of *Satyrichthys kikingeri* \(Teleostei: Peristediidae\) from the Maldives Archipelago \(Indian Ocean\)](#). Marine Biodiversity Records 7(e61): 1–4.
- Porter MM, Novitskaya E, Castro-Ceseña AB, Meyers MA, McKittrick J (2013) [Highly deformable bones: unusual deformation mechanisms of seahorse armor](#). Acta Biomaterialia 9: 6763–6770.
- Qu Q, Zhu M, Wang W (2013) [Scales and dermal skeletal histology of an early bony fish *Psarolepis romeri* and their bearing on the evolution of rhombic scales and hard tissues](#). PLoS One 8: e61485.
- Van Oijen MJ, Kawai T, Loots I (2013) [Putative type specimens of *Satyrichthys* \(Scorpaeniformes: Peristediidae\) in the Bleeker collection of the Naturalis Biodiversity Center, Leiden, The Netherlands](#). Zootaxa 3670: 207–

214.

Yang W, Gludovatz B, Zimmermann EA, Bale HA, Ritchie RO, Meyers MA (2013) [Structure and fracture resistance of alligator gar \(*Atractosteus spatula*\) armored fish scales](#). Acta Biomaterialia 9: 5876–5889.

Yang W, Naleway SE, Porter MM, Meyers MA, McKittrick J (2015) The armored carapace of the boxfish. Acta Biomaterialia 23: 1–10.



A Ngamniyom  <https://orcid.org/0000-0002-2325-0622>