



## Study on proximate composition of four seaweeds from Kilinochchi and Kalpitiya area of Sri Lanka

Vidushika Sanjeevani Premadasa • Dissanayake Mudiyansele Anusha Edirisinghe

Department of Aquaculture and Aquatic Resources Management, University College of Anuradhapura, University of Vocational Technology, Sri Lanka

### Correspondence

Dissanayake Mudiyansele Anusha Edirisinghe; Department of Aquaculture and Aquatic Resources Management, University College of Anuradhapura, University of Vocational Technology, Sri Lanka

 [anushaedirisinghe@gmail.com](mailto:anushaedirisinghe@gmail.com)

### Manuscript history

Received 13 October 2020 | Accepted 27 May 2022 | Published online 7 June 2022

### Citation

Premadasa VS, Edirisinghe DMA (2022) Study on proximate composition of four seaweeds from Kilinochchi and Kalpitiya area of Sri Lanka. *Journal of Fisheries* 10(2): 102401. DOI: 10.17017/j.fish.286

### Abstract

Some of the representatives from edible seaweeds (*Gracilaria verrucosa*, *Gracilaria corticata*, *Turbinaria ornata*, *Sargassum polycystum*) from Kilinochchi and Kalpitiya in Sri Lanka were investigated for proximate composition (moisture, protein, lipid and ash content). The proximate composition was significantly ( $p < 0.001$ ) different across seaweed species. In this regard, moisture content (mean  $\pm$  SD) varied from  $76.33 \pm 0.0\%$  in *T. ornata* to  $89.78 \pm 0.6\%$  in *G. corticata*. The *S. polycystum* presented the highest fat content ( $1.94 \pm 0.5\%$ ); whereas, *G. corticata* showed the highest protein level ( $21.66 \pm 1.1\%$ ). Fat content was low in *T. ornata* ( $0.37 \pm 0.2\%$ ) and *S. polycystum* ( $1.94 \pm 0.5\%$ ) while protein content was low in other three species. Also *G. verrucosa* and *G. corticata* had higher ash content ( $44.34 \pm 3.8\%$  and  $44.34 \pm 3.8\%$  respectively) than other seaweed species. The variation in proximate composition was high among species and would be a healthy addition to normal diets.

**Keywords:** ash; lipid; moisture; protein; seaweeds

### 1 | INTRODUCTION

The Sri Lankan seaweed flora is relatively rich, with about 440 taxa, belonging to 148 genera currently recorded along a coastline of 1585 km (Barton 1903; Boergesen 1936; Durairatnam 1961; Baldwin 1991). They are plant like organisms that generally live attached to the rock or hard substrata in the coastal areas. There is ample evidence of the importance of seaweeds in the context of human food needs and ecological benefits of farming (Craigie 2011; Cornish *et al.* 2017). Economic uses of seaweeds particularly for food, feed, phycocolloids and agro-based products, are well known for centuries and are now being extensively investigated for their application in bio-fuel, nutraceutical, medicinal, personal-care and food additive industries (Hafting *et al.* 2015; Kim *et al.* 2017).

There are three major classes of seaweed namely, Chlorophyceae (green), Rhodophyceae (Red) and Phaeophyceae (Brown) (Trono Jr and Ganzon Ganzon-Fortes 1988). The red, brown and green seaweeds are phylogenetically very different and this is reflected in their differences in growth, structure, and biochemical composition (Lange *et al.* 2020). The most common seaweeds found in Sri Lanka are the brown seaweeds belonging to the genus *Sargassum* (Jayasuriya *et al.* 1990). In this study, *Gracilaria verrucosa* and *Gracilaria corticata* of Rhodophyta and *Turbinaria ornata* and *Sargassum polycystum* of Phaeophyceae were studied for proximate composition.

Seaweeds are not used as much for edible use although many kinds of seaweeds are collected in Sri Lanka

(Komatsuzaki *et al.* 2019). Since 1800s, naturally collected seaweed species have been exported from Sri Lanka (Duraitarnam 1963) and a growing export market was identified thereafter. Furthermore, a small percentage of dried seaweeds is being sold locally, while a good demand for packed *Gracilaria* spp. can be identified in Islamic festive season in Sri Lanka (Jayasuriya 1987). *Gracilaria* spp. are popular food item among fishermen, especially in producing areas and was domestically consumed as porridge and a jelly drink (Ginigaddara *et al.* 2018).

Different species of seaweeds especially protein rich seaweeds are used as human food in many countries all over the world. Therefore, studies on the proximate composition of seaweeds are important to determine their nutritive value (Dawes 1981).

## 2 | METHODOLOGY

The four seaweed species namely, *G. verrucosa*, *G. corticata*, *T. ornate* and *S. polycystum* were collected from Kalpitiya and Kilinochchi coastal regions of Sri Lanka. Seaweed samples were picked with hand from natural beds and immediately washed with seawater to remove the foreign particles, sand particles and epiphytes. Then they were kept in air tight bags in an ice box containing ice flakes and immediately transported to the laboratory in the department of Aquaculture and Aquatic Resources Management, University College of Anuradhapura. The samples were washed thoroughly using running water to remove the salt on the surface of the sample. The samples were dried in oven at 40°C to constant weight. Then dried samples were grounded in an electric mixer (Trio – MG 158, India) and powdered samples were stored in Chest freezer (HCF775, China) until further analysis. Protein, moisture and ash content of the seaweeds were determined using the AOAC standard methods with triplicate samples.

The moisture content was determined by using the moisture meter at 105°C to constant weight (AOAC 1989). The crude protein content was analysed by the semiautomated Kjeldahl system (Nitrogen distiller; Raypa DNP-

2000MP, Spain), which was following the AOAC official method of analysis 984.13 (AOAC 2005). Ash content was analysed by incinerating the seaweeds in the muffle furnace (LabTech LEF-P, Korea) at 550°C for 4 hours by following the AOAC official method of analysis (AOAC 1989). The total lipid content was evaluated using Bligh and Dyer (1959) method.

Data were analysed with SPSS 16.0 software using parametric tests. One-way ANOVA was used to compare the means of chemical composition data determined for the four seaweed species. When significant differences were found, Duncan multiple comparison test was used, and differences were considered significant at  $p < 0.05$ .

## 3 | RESULTS AND DISCUSSION

Chemical compositions were significantly ( $p < 0.05$ ) different among the four seaweed species. The moisture content of fresh seaweed samples were ranged from 76.33% to 89.78% (Table 1). Among the four seaweed studied, *G. corticata* ( $89.78 \pm 0.6\%$ ) was found to have the highest moisture content followed by *S. polycystum* (mean  $\pm$  SD:  $80.25 \pm 0.4\%$ ), *G. verrucosa* ( $79.78 \pm 0.7\%$ ) and *T. ornate* ( $76.33 \pm 0.0\%$ ). The moisture content of four seaweeds was close to the value (79 – 86%) previously reported by Jayasinghe *et al.* (2018).

The ash content in most seaweed varieties is much higher than terrestrial plants (Jayasinghe *et al.* 2018). The ash content of studied seaweeds was between  $26.11 \pm 0.9\%$  and  $46.72 \pm 0.5\%$ . The amount of ash content obtained in the present study was in agreement with previous studies which stated that seaweeds contained high proportions of ash (21.1 – 39.3%) (Rupérez 2002; Mohammadi *et al.* 2013.). During this research, it was observed that, the red algae (*G. verrucosa* and *G. corticata*) had significantly higher ash (46.72 – 44.34%) than the brown algae (*T. ornate*, 26.11%; *S. polycystum*, 29.45%). The deference of the ash content depends on the seaweed species, a method of mineralisation, types of processing, marine environmental factors and physiological feature of the seaweed (Rameshkumar *et al.* 2013).

**TABLE 1** Proximate composition (moisture, ash, lipid and protein) of selected seaweeds in % dry weight.

| Seaweed species             | Moisture (%)      | Ash (%)           | Lipid (%)        | Protein (%)       |
|-----------------------------|-------------------|-------------------|------------------|-------------------|
| <i>Gracilaria verrucosa</i> | $79.78 \pm 0.7^b$ | $46.72 \pm 0.5^a$ | $0.20 \pm 0.0^b$ | $16.51 \pm 1.7^b$ |
| <i>Gracilaria corticata</i> | $89.78 \pm 0.6^a$ | $44.34 \pm 3.8^a$ | $0.24 \pm 0.1^b$ | $21.66 \pm 1.1^a$ |
| <i>Turbinaria ornata</i>    | $76.33 \pm 0.0^c$ | $26.11 \pm 0.9^b$ | $0.37 \pm 0.2^b$ | $16.79 \pm 0.2^b$ |
| <i>Sargassum polycystum</i> | $80.25 \pm 0.4^b$ | $29.45 \pm 0.7^b$ | $1.94 \pm 0.5^a$ | $17.80 \pm 0.1^b$ |

Values are means of triplicate groups  $\pm$  SD,  $n = 3$ .

Seaweeds are relatively low in lipid (1 – 5% of dry weight) (Burtin 2003; Polat and Ozogul 2008). In the present study, the highest crude lipid content was observed in *S. polycystum* (1.94 %) followed by *T. ornata* (0.37 %), *G. corticata* (0.24%) and *G. verrucosa* (0.20%). These results are comparably lower than those reported by Cirik

*et al.* (2010); Rohani-Ghadikolaei *et al.* (2012); Remya *et al.* (2019) and Balboa *et al.* (2016). But this research is in accordance with Jayasinghe *et al.* (2018) who reported that the total lipid content was generally low in all seaweeds samples (0.30% to 1.97% in dry weight) and form a minor portion of the proximate component than other

components. These differences in results between studies could be due to the different environmental conditions, the season of harvesting and the habitat of the seaweeds.

The protein content is one of the important constituents and suggested as supplemental food for humans and animals (Jayasinghe *et al.* 2018). In this study, the highest protein level was found in *G. corticata* (21.66%) followed by *S. polycystum* (17.8%), *T. ornata* (16.79%) and *G. verrucosa* (16.51%) (Table 1). These levels of the protein content of the seaweeds depend on the environmental factors, maturity stage, seasonality and the physical feature of the seaweeds plants (Banerjee *et al.* 2009)

#### 4 | CONCLUSIONS

From the present study, it is obvious that marine macro algae are rich in nutritive properties. The present findings could be useful to collect the selected seaweeds from Northern and North Western coast of Sri Lanka and use them in the food and pharmaceutical industries for various purposes. The results of the study concluded that these seaweeds can provide dietary alternatives to improve the nutritive value of the human diet. Further study needs to be done on the utilisation and sensory insights of these seaweeds.

#### CONFLICT OF INTEREST

The authors declare no conflict of interest.

#### AUTHORS' CONTRIBUTION

**VSP** Data collection and sample analysis; **DMAE** data analysis, manuscript preparation and research supervision.

#### DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available on request from the corresponding author.

#### REFERENCES

- AOAC (1989) Official methods of analysis 15th edition. Association of Official Analytical Chemists, Washington DC.
- AOAC (2005) Method number 984.13. Official methods of analysis of AOAC International. Association of Official Analytical Chemists, Washington DC.
- Balboa EM, Gallego-Fábrega C, Moure A, Domínguez H (2016) Study of the seasonal variation on proximate composition of oven-dried *Sargassum muticum* biomass collected in Vigo Ria, Spain. *Journal of Applied Phycology* 28(3): 1943–1953.
- Baldwin MF (Ed) (1991) Natural resources of Sri Lanka: conditions and trends. Keels Business Systems Ltd. Education Center, Sri Lanka.
- Banerjee K, Ghosh R, Homechaudhuri S, Mitra A (2009) Seasonal variation in the biochemical composition of red seaweed (*Catenella repens*) from Gangetic Delta, NorthEast coast of India. *Journal of Earth System Science* 118(5): 497–505.
- Barton ES (1903) List of marine algae collected by Prof. Herdman at Ceylon in 1902, with a note on the fructification of *Halimeda*. Herdman Ref. on Ceylon Pearl Fishery Part 1: 168–169.
- Bligh EG, Dyer WJ (1959) A rapid method of total lipid extraction and purification. *Canadian Journal of Biochemistry and Physiology* 37: 911–917.
- Boergensen F (1936) Some marine algae from Ceylon. *Ceylon Journal of Science* 12 2: 57–96.
- Burtin P (2003) Nutritional value of seaweeds. *Electronic Journal of Environmental Agricultural and Food Chemistry* 2: 498–503.
- Cirik Ş, Çetin Z, Ak İ, Cirik S, Göksan T (2010) Greenhouse cultivation of *Gracilaria verrucosa* (Hudson) papenfuss and determination of chemical composition. *Turkish Journal of Fisheries and Aquatic Sciences* 10(4): 559–564.
- Cornish M, Critchley AT, Mouritsen O (2017) Consumption of seaweeds and the human brain. *Journal of Applied Phycology* 29: 2377–2398.
- Craigie JSJ (2011) Seaweed extract stimuli in plant science and agriculture. *Journal of Applied Phycology* 23: 371–393.
- Dawes CJ (1981) Marine botany. John Wiley and sons Inc, New York. 508 pp.
- Durairatnam M (1961) Contribution to the study of marine algae of Ceylon. *Bulletin of the Fisheries Research Station Sri Lanka (Ceylon)* 10: 5–117.
- Durairatnam M (1963) Some marine algae from Ceylon 2 (*Laurencia lamourous*). *Bulletin of the Fisheries Research Station* 16(2): 19–28.
- Ginigaddara GAS, Lankapura AIY, Rupasena LP, Bandara AMKR (2018) Seaweed farming as a sustainable livelihood option for northern coastal communities in Sri Lanka. *Future of Food: Journal on Food, Agriculture and Society* 6(1): 57–70.
- Hafting JJ, Craigie JS, Stengel DB, Loureiro RR, Buschmann AH, ... Critchley AT (2015) [Prospects and challenges for industrial production of seaweed bioactives](#). *Journal of Phycology* 51: 821–837.
- Jayasinghe GDTM, Jinadasa BKKK, Chinthaka SDM (2018) Nutritional composition and heavy metal content of five tropical seaweeds. *The Open Analytical Chemistry Journal* 3(2): 17–22.
- Jayasuriya PMA (1987) The socio economic aspects of the people traditionally involved in the harvesting of wild seaweeds (*Gracilaria lichenoides*) from Puttalam lagoon. *Proceedings of 42nd Annual sessions of Sri Lanka Association for the Advancement of Science*.
- Jayasuriya PMA, Island C, Colombo M (1990) The status of culture and utilization of seaweeds in Sri Lanka. In: *Report of the Regional Workshop on the Culture &*

Utilization of Seaweeds. FAO Corporate Document Repository.

- Kim JK, Yarish C, Hwang EK, Park M, Kim Y (2017) [Seaweed aquaculture: cultivation technologies, challenges and its ecosystem services](#). *Algae* 32: 1–13.
- Komatsuzaki N, Arai S, Fujihara S, Shima J, Wijesekara RS, de Croos MDS (2019) [Development of novel bread by combining seaweed \*Kappaphycus alvarezii\* from Sri Lanka and \*Saccharomyces cerevisiae\* isolated from Nectarine](#). *Journal of Agricultural Science and Technology B* 9: 339–346.
- Lange L, Bak UG, Hansen SCB, Gregersen O, Harmsen P, ... Hreggviðsson GÓ (2020) [Opportunities for seaweed biorefinery](#). In: *Sustainable Seaweed Technologies Cultivation, Biorefinery, and Applications*. *Advances in Green and Sustainable Chemistry*. Elsevier: 3-31.
- Mohammadi M, Tajik H, Hajeb P (2013) Nutritional composition of seaweeds from the Northern Persian Gulf. *Iranian Journal of Fisheries Sciences* 12(1): 232–240.
- Polat S, Ozogul Y (2008) Biochemical composition of some red and brown macro-algae from the Northeastern Mediterranean Sea. *International Journal of Food Science and Nutrition* 59: 566–572.
- Rameshkumar S, Ramakritinan CM, Yokeshbabu M (2013) Proximate composition of some selected seaweeds from Palk Bay and Gulf of Mannar, Tamilnadu, India. *Asian Journal of Biomedical and Pharmaceutical Sciences* 3(16): 1.
- Remya RR, Rajasree SR, Suman TY, Aranganathan L, Gayathri S, ... Karthih MG (2019) [Studies on proximate composition and phytochemical profiling of \*Turbinaria ornata\* and its anti-proliferative effect on Y79 cell lines](#). *Thalassas: An International Journal of Marine Sciences* 35(2): 495–502.
- Rohani-Ghadikolaei K, Abdulalian E, Ng W (2012) [Evaluation of the proximate, fatty acid and mineral composition of representative green, brown and red seaweeds from the Persian Gulf of Iran as potential food and feed resources](#). *Journal of Food Science and Technology* 49: 774–780.
- Rupérez P (2002) Mineral content of edible marine seaweeds. *Food Chemistry* 79(1): 23–26.
- Trono Jr GC, Ganzon-Fortes ET (1988) Philippines seaweeds. Technology and Livelihood Center PCPM Certificate of Registration. No. 594.



**VS Premadasa**  <https://orcid.org/0000-0001-9314-6540>

**DMA Edirisinghe**  <https://orcid.org/0000-0001-7155-6696>