

Effects of *Streptococcus agalactiae* infection and oral florfenicol administration on the hemato-biochemistry, erythrocyte morphology and histopathology of *Oreochromis niloticus*

Joshi Sharon¹ • Thangapalam Jawahar Abraham¹ • Arya Sen¹ • Ratnapriya Das¹ • Priyanka Sinha¹ • Satyanarayana Boda² • Arumugam Uma³ • Prasanna Kumar Patil⁴

¹ Department of Aquatic Animal Health, Faculty of Fishery Sciences, West Bengal University of Animal and Fishery Sciences, Chakgaria, Kolkata 700094, West Bengal, India

² Department of Fishery Economics and Statistics, Faculty of Fishery Sciences, West Bengal University of Animal and Fishery Sciences, Chakgaria, Kolkata 700094, West Bengal, India

³ State Referral Laboratory for Aquatic Animal Health, Tamil Nadu Dr J.Jayalalithaa Fisheries University, Madhavaram Milk Colony, Chennai 600051, Tamil Nadu, India

⁴ Aquatic Animal Health and Environment Division, ICAR-Central Institute of Brackishwater Aquaculture, Raja Annamalai Puram, Chennai 600028, Tamil Nadu, India

Article link: <https://doi.org/10.17017/j.fish.720>

Citation:

Sharon J, Abraham TJ, Sen A, Das R, Sinha P, Boda S, Uma A, Patil PK (2025) Effects of *Streptococcus agalactiae* infection and oral florfenicol administration on the hemato-biochemistry, erythrocyte morphology and histopathology of *Oreochromis niloticus*. Journal of Fisheries 13(1): 131209. DOI: 10.17017/j.fish.720




Figure S1 Gross and clinical changes, wound progression and healing, and discoloration observed in *Streptococcus agalactiae* LCR1 challenged and florfenicol (FFC) fed *Oreochromis niloticus* juveniles. Florfenicol treatment: Fish were challenged with *S. agalactiae* LCR1 and fed FFC at $15 \text{ mg kg biomass}^{-1} \text{ day}^{-1}$ for 10 consecutive days. Untreated: Fish were challenged with *S. agalactiae* LCR1 and fed a control diet. DPI: Day post injection.