



Ichthyofaunal diversity and fishery status of Sutrapada Coast, Gujarat, India


Santosh Kumar Singh¹ • Kangkan Jyoti Sarma² • Dhaval M. Bhatt¹ • Pradeep C. Mankodi¹

¹ Division of Marine and Freshwater Biology, Department of Zoology, Faculty of Science, The Maharaja Sayajirao University of Baroda, Vadodara – 390020, India

² Division of Fish and Fishery Biology, Department of Zoology, School of Biological Sciences, University of Science & Technology Meghalaya, Ri-Bhoi, Meghalaya-793101, India

Correspondence

Santosh Kumar Singh; Division of Marine and Freshwater Biology, Department of Zoology, Faculty of Science, The Maharaja Sayajirao University of Baroda, Vadodara – 390020, India

 santosh123.14689@gmail.com

Manuscript history

Received 12 November 2020 | Revised 5 June 2021 | Accepted 12 June 2021 | Published online 12 July 2021

Citation

Singh SK, Sarma KJ, Bhatt DM, Mankodi PC (2021) Ichthyofaunal diversity and fishery status of Sutrapada Coast, Gujarat, India. *Journal of Fisheries* 9(2): 92204.

Abstract

Gujarat being a state having largest coastline and highest continental shelf in India represents diverse coastal habitats. Gujarat has vast potential of marine bioresources and offer considerable scope for marine fisheries. The ichthyofaunal diversity of Sutrapada, Gir Somnath district, Gujarat was investigated for a period of three years from June 2015 to June 2018 through the analysis of fish catch data of trawl fisheries and achieves of Fisheries Department, Central Institute of Fisheries Technology - Central Marine Fisheries Research Institute (CIFT-CMFRI), Marine Products Export Development Authority (MPEDA) and Commissioner Fisheries Office, Veraval. During this study period, 114 fish species belonging to 55 families and 93 genera were identified. The paper emphasizes the importance of further research on respective families for understanding of its distribution, ecology and conservation status.

Keywords: Diversity; Sutrapada; fish; species composition

1 | INTRODUCTION

Fishes are the largest group of vertebrates, which exhibit a remarkable diversity of morphological attributes and biological adaptations. Species identification is challenging for taxonomists (Zhang and Hanner 2011) when facing new biota. Globally, ichthyofaunal diversity comprises approximately half of the total members of the subphylum Vertebrata with 35588 valid fish species (Fricke *et al.* 2020). In India, ichthyofauna from the fresh and marine waters account for 9.7 percent of the global population, among which the marine fishes alone account for 7.4 percent (Eschmeyer and Fong 2014). Day (1899a, 1899b) reported 1418 species of fish belonging to 342 genera from British India whereas 2546 species of ichthyofauna from 969 genera, 254 families and 40 orders were de-

scribed by Talwar (1991). Recent findings have increased number of valid fish species in India with an estimation of 3231 species of freshwater, brackish water and marine species (Gopi and Mishra 2015). Out of the total fish diversity, marine water supports 2443 species whereas freshwater supports 675 species (Gopi and Mishra 2015) and brackish water comprises approximately 113 species (Sarkar *et al.* 2012). Fishes are a major source of human diet (Sarwade and Khillare 2010; Galib *et al.* 2013) as well as of oil, fertilizer and feed for domestic animals.

The state of Gujarat coastline is about 20% of Indian coastline, 33% of the continental shelf area (164000 square km) and over 200000 square km of Exclusive Economic Zone (EEZ) and ranks first among India's nine maritime states in marine fish production (Mohanraj *et al.*

2009). Saurashtra Coast represents major portion of the Gujarat coastline, having 26 landing centers which is 78% of the total landings of the state (Balan *et al.* 1987). Saurashtra coastline – peninsula of Gujarat is having major fish landing centres such as Okha, Porbandar, Mangrol, Veraval, Sutrapada, Navabandar, Jafarabad etc. Gujarat is the leading state in marine fish production since 2012 (Frad, CMFRI 2018). In 2016, the total landing of the marine fishes was 0.77 metric tonnes, which constitutes around 21.32% to the total fish landings of India. In Gujarat state, Gir-Somnath district contributes maximum landings (0.34 lakh tonnes) which is approximately 44% of the total landings of Gujarat (Joshi *et al.* 2018). Sutrapada is the second major landing centre of the Gir-Somnath district, Gujarat. To date, there are no studies carried out on the diversity of the ichthyofauna from Sutrapada Coast but study of ichthyofauna and their status are essential for the sustainable management of the fishery (Galib 2015; Galib *et al.* 2009, 2016; Mohsin *et al.* 2009, 2013). In Sutrapada, the fishery constitutes primarily of pomfrets, seer fishes, croakers, hilsa and other clupeoids, catfish, ribbonfish, perches and silver bass. The aim of this study was to investigate the current ichthyofaunal diversity of Sutrapada. The study outcomes will be helpful in the management and conservation of the commercially and ecologically important fishes.

2 | METHODOLOGY

In Sutrapada village - bander, about 381 Out Boat Motors (OBM) gillnetters are being operated of which most are involved in fishing. These vessels are made up of Fibre Reinforced Plastics (FRP). In this study data were collected from 23 OBM gillnetters. Fishermen in the study area usually go for 2 – 3 days fishing trip but one day trip is also common. Multifilament and monofilament nylon

nets operated by FRP canoes fitted with outboard machine were being used for fishing.

The fish specimens were collected with the help of professional fishermen from Sutrapada Coast (20°50'11.94"N, 70°28'52.48"E) from June 2015 to June 2019. The collection of the fishes was performed on a monthly basis during winter (November to February) and summer (March to June) seasons. In monsoon (July to October) season fishing activities are closed due to safety reasons thus, no collection was made during this period. After collection, on field photographs were taken and morphological characters were recorded. The meristic and morphometric characters of collected specimens were measured and used for identification of the species following standard process and literature (Day 1878; Jhingran 1983; Sarwade and Khillare 2010; Murugan and Namboothri 2012). Conservation categories of the fish species are based on Redlist database of International Union for Conservation of Nature and Natural Resources (IUCN 2021).

3 | RESULT AND DISCUSSION

The marine fishery recourses of Sutrapada consisted exclusively of capture fisheries. Many collected fishes are having economic importance and sold after collection in the local fish market (Sarwade and Khillare 2010). During this study period, 114 fish species belonging to two classes, 18 orders, 55 families and 93 genera were collected and identified (Table 1). The most dominant family found was Carangidae ($n = 15$ species) followed by Scombridae ($n = 9$), Sciaenidae ($n = 6$), Carcharhinidae ($n = 4$), Synodontidae ($n = 4$), Ariidae ($n = 4$), Clupeidae ($n = 3$), Engraulidae ($n = 3$), Nemipteridae ($n = 3$) and rest families constituted of single or double species (Table 1).

TABLE 1 Fish species collected from Sutrapada Coast. Status of the species is based on IUCN (2021). EN, Endangered; DD, Data Deficient; LC, Least Concern; NE, Not Evaluated; NT, Near Threatened; VU, Vulnerable.

Class/ Order/ Family	Species	English name	Status
Class: Elasmobranchii			
Order: Carcharhiniformes			
Carcharhinidae	<i>Carcharhinus limbatus</i> (Müller & Henle, 1839)	Blacktip shark	VU
	<i>Scoliodon laticaudus</i> Müller & Henle 1838	Spadenose shark	NT
	<i>Carcharhinus dussumieri</i> (Müller & Henle 1839)	Whitecheek shark	EN
	<i>Carcharhinus sorrah</i> (Müller & Henle 1839)	Spot-tail shark	NT
Triakidae	<i>Mustelus mosis</i> Hemprich & Ehrenberg, 1899	Arabian smooth-hound	DD
Order: Myliobatiformes			
Dasyatidae	<i>Maculabatis gerrardi</i> (Gray, 1851)	Sharpnose stingray	VU
Order: Rhinopristiformes			
Rhinidae	<i>Rhynchobatus djiddensis</i> (Forsskål 1775)	Giant guitarfish	CR
Rhinobatidae	<i>Rhinobatos punctifer</i> Compagno & Randall, 1987	Spotted guitarfish	NT
Glaucostegidae	<i>Glaucostegus granulatus</i> (Cuvier 1829)	Granulated guitarfish	CR

TABLE 1 Continued.

Class/ Order/ Family	Species	English name	Status	
Order: Torpediniformes				
Torpedinidae	<i>Torpedo fuscomaculata</i> Peters, 1855	Black-spotted torpedo	DD	
	<i>Torpedo sinuspersici</i> Olfers, 1831	Variable torpedo ray	DD	
Class: Actinopterygii				
Order: Acanthuriformes				
Acanthuridae	<i>Acanthurus mata</i> (Cuvier, 1829)	Elongate surgeonfish	NE	
	<i>Zebрасoma desjardini</i> (Bennett, 1836)	Indian sail-fin surgeonfish	LC	
Chaetodontidae	<i>Chaetodon collare</i> Bloch, 1787	Redtail butterflyfish	LC	
Pomacanthidae	<i>Pomacanthus annularis</i> (Bloch, 1787)	Bluering angelfish	LC	
Ephippidae	<i>Platax teira</i> (Forsskål, 1775)	Longfin batfish	NE	
Order: Anguilliformes				
Muraenesocidae	<i>Congresox talabonoides</i> (Bleeker 1853)	Indian pike conger	NE	
	<i>Muraenesox cinereus</i> (Forsskål 1775)	Daggertooth pike conger	LC	
Order: Aulopiformes				
Synodontidae	<i>Saurida tumbil</i> (Bloch, 1795)	Greater lizardfish	LC	
	<i>Saurida longimanus</i> Norman 1939	Longfin lizardfish	LC	
	<i>Trachinocephalus myops</i> (Forster, 1801)	Snakefish lizard fish	LC	
	<i>Harpadon nehereus</i> (Hamilton 1822)	Bombay-duck	NT	
Order: Beloniformes				
Belonidae	<i>Tylosurus crocodilus</i> (Péron & Lesueur 1821)	Hound needlefish	LC	
	<i>Strongylura strongyle</i> (van Hasselt 1823)	Spottail needlefish	LC	
Exocoetidae	<i>Parexocoetus brachypterus</i> (Richardson 1846)	Sailfin flyingfish	DD	
	<i>Hirundichthys coromandelensis</i> (Hornell 1923)	Coromandel flyingfish	LC	
Hemiramphidae	<i>Hemiramphus far</i> (Forsskål 1775)	Black-barred halfbeak	NE	
Order: Carangiformes				
Carangidae	<i>Atropus atropos</i> (Bloch & Schneider, 1801)	Cleftbelly trevally	LC	
	<i>Decapterus russelli</i> (Rüppell, 1830)	Indian scad	LC	
	<i>Decapterus macrosoma</i> Bleeker 1851	Shortfin scad	LC	
	<i>Megalaspis cordyla</i> (Linnaeus 1758)	Torpedo scad	LC	
	<i>Alectis indica</i> (Rüppell, 1830)	Indian threadfish	LC	
	<i>Alepes kleinii</i> (Bloch, 1793)	Razorbelly scad	LC	
	<i>Elagatis bipinnulata</i> (Quoy & Gaimard, 1825)	Rainbow runner	LC	
	<i>Parastromateus niger</i> (Bloch, 1795)	Black pomfret	LC	
	<i>Scomberoides tol</i> (Cuvier, 1832)	Needlescaled queenfish	LC	
	<i>Scomberoides commersonianus</i> Lacepède, 1801	Talang queenfish	LC	
	<i>Alepes djedaba</i> (Forsskål 1775)	Shrimp scad	LC	
	<i>Caranx sexfasciatus</i> Quoy & Gaimard 1825	Bigeye trevally	LC	
	<i>Carangoides malabaricus</i> (Bloch & Schneider 1801)	Malabar trevally	LC	
	<i>Carangoides coeruleopinnatus</i> (Rüppell 1830)	Coastal trevally	LC	
	<i>Atule mate</i> (Cuvier 1833)	Yellowtail scad	LC	
	Sphyraenidae	<i>Sphyraena jello</i> Cuvier 1829	Pickhandle barracuda	LC
		<i>Sphyraena putnamae</i> Jordan & Seale 1905	Sawtooth barracuda	DD
	Polynemidae	<i>Eleutheronema tetradactylum</i> (Shaw 1804)	Fourfinger threadfin	EN
		<i>Leptomelanosoma indicum</i> (Shaw 1804)	Indian threadfin	LC
Istiophoridae	<i>Istiompax indica</i> (Cuvier 1832)	Black marlin	DD	
	<i>Istiophorus platypterus</i> (Shaw 1792)	Indo-Pacific sailfish	LC	
Coryphaenidae	<i>Coryphaena hippurus</i> Linnaeus, 1758	Common dolphinfish	LC	
Echeneidae	<i>Echeneis naucrates</i> Linnaeus, 1758	Live sharksucker	LC	
Menidae	<i>Mene maculata</i> (Bloch & Schneider, 1801)	Moonfish	NE	
Polynemidae	<i>Leptomelanosoma indicum</i> (Shaw, 1804)	Indian threadfin	NE	
Rachycentridae	<i>Rachycentron canadum</i> (Linnaeus, 1766)	Cobia	LC	
Xiphiidae	<i>Xiphias gladius</i> Linnaeus 1758	Swordfish	LC	

TABLE 1 Continued.

Class/ Order/ Family	Species	English name	Status
Order: Centrarchiformes			
Kyphosidae	<i>Kyphosus vaigiensis</i> (Quoy & Gaimard, 1825)	Brassy chub	LC
Terapontidae	<i>Terapon jarbua</i> (Forsskål, 1775)	Jarbua terapon	LC
Order: Clupeiformes			
Clupeidae	<i>Sardinella gibbosa</i> (Bleeker, 1849)	Goldstripe sardinella	LC
	<i>Sardinella longiceps</i> Valenciennes, 1847	Indian oil sardine	LC
	<i>Tenualosa ilisha</i> (Hamilton 1822)	Hilsa shad	LC
Chirocentridae	<i>Chirocentrus nudus</i> Swainson 1839	Whitefin wolf-herring	LC
	<i>Chirocentrus dorab</i> (Forsskål 1775)	Dorab wolf-herring	LC
Dussumeriidae	<i>Dussumieria acuta</i> Valenciennes, 1847	Rainbow sardine	LC
Engraulidae	<i>Thryssa setirostris</i> (Broussonet, 1782)	Longjaw thryssa	LC
	<i>Thryssa dussumieri</i> (Valenciennes, 1848)	Dussumier's thryssa	LC
	<i>Thryssa malabarica</i> (Bloch, 1795)	Malabar thryssa	DD
Pristigasteridae	<i>Ilisha megaloptera</i> (Swainson 1839)	Bigeye ilisha	LC
	<i>Opisthopterus tardoore</i> (Cuvier 1829)	Tardoore	LC
Order: Mugiliformes			
Mugilidae	<i>Mugil cephalus</i> Linnaeus, 1758	Grey mullet	LC
Order: Perciformes			
Haemulidae	<i>Pomadasys maculatus</i> (Bloch, 1793)	Saddle grunt	LC
	<i>Pomadasys argenteus</i> (Forsskål 1775)	Silver grunt	LC
Priacanthidae	<i>Priacanthus hamrur</i> (Forsskål, 1775)	Moontail bullseye	LC
Serranidae	<i>Cephalopholis sonnerati</i> (Valenciennes, 1828)	Tomato hind	LC
	<i>Epinephelus diacanthus</i> (Valenciennes, 1828)	Spinycheek grouper	LC
Sillaginidae	<i>Sillago sihama</i> (Forsskål, 1775)	Silver sillago	LC
Lethrinidae	<i>Lethrinus ornatus</i> Valenciennes 1830	Ornate emperor	LC
	<i>Lethrinus nebulosus</i> (Forsskål, 1775)	Spangled emperor	LC
Lutjanidae	<i>Lutjanus johnii</i> (Bloch, 1792)	John's snapper	LC
Sparidae	<i>Argyrops spinifer</i> (Forsskål, 1775)	King soldier bream	LC
Uranoscopidae	<i>Uranoscopus archionema</i> Regan, 1921	–	NE
Nemipteridae	<i>Nemipterus japonicus</i> (Bloch, 1791)	Japanese threadfin bream	NE
	<i>Scolopsis vosmeri</i> (Bloch, 1792)	Whitecheek monocle bream	NE
	<i>Parasclopsis eriomma</i> (Jordan & Richardson, 1909)	Swallowtail dwarf monocle bream	NE
Platycephalidae	<i>Platycephalus indicus</i> (Linnaeus, 1758)	Bartail flathead	DD
Sciaenidae	<i>Otolithoides biauritus</i> (Cantor, 1849)	Bronze croaker	NE
	<i>Nibea maculata</i> (Bloch & Schneider 1801)	Blotched croaker	NE
	<i>Otolithes cuvieri</i> Trewavas 1974	Lesser tigertooth croaker	LC
	<i>Johnius dussumieri</i> (Cuvier 1830)	Sin croaker/ Bearded croaker	LC
	<i>Otolithes ruber</i> (Bloch & Schneider 1801)	Tiger toothed croaker	LC
	<i>Protonibea diacanthus</i> (Lacepède 1802)	Blackspotted croaker/Ghol	LC
Scorpaenidae	<i>Pterois miles</i> (Bennett, 1828)	Devil firefish	LC
Order: Pleuronectiformes			
Paralichthyidae	<i>Pseudorhombus arsius</i> (Hamilton 1822)	Largetooth flounder	LC
Psettodidae	<i>Psettodes erumei</i> (Bloch & Schneider 1801)	Indian halibut	DD
Order: Scombriformes			
Scombridae	<i>Auxis rochei</i> (Risso, 1810)	Bullet tuna	LC
	<i>Euthynnus affinis</i> (Cantor 1849)	Little tuna	LC
	<i>Auxis thazard</i> (Lacepède 1800)	Frigate tuna	LC
	<i>Rastrelliger kanagurta</i> (Cuvier 1816)	Indian mackerel	DD
	<i>Katsuwonus pelamis</i> (Linnaeus 1758)	Skipjack tuna	LC
	<i>Thunnus albacares</i> (Bonnaterre 1788)	Yellow fin tuna	NT
	<i>Thunnus tonggol</i> (Bleeker 1851)	Longtail tuna	DD

TABLE 1 Continued.

Class/ Order/ Family	Species	English name	Status
Scombridae	<i>Thunnus obesus</i> (Lowe 1839)	Bigeye tuna	VU
	<i>Scomberomorus guttatus</i> (Bloch & Schneider 1801)	Indo-Pacific king mackerel	DD
Stromateidae	<i>Pampus argenteus</i> (Euphrasen, 1788)	Silver pomfret	VU
	<i>Pampus chinensis</i> (Euphrasen 1788)	Chinese silver pomfret	DD
Trichiuridae	<i>Trichiurus lepturus</i> Linnaeus, 1758	Largehead hairtail fish	LC
	<i>Lepturacanthus savala</i> (Cuvier 1829)	Savalai hairtail/Ribbon fish	DD
Order: Siluriformes			
Ariidae	<i>Plicofollis layardi</i> (Günther, 1866)	Thinspine sea catfish	NE
	<i>Plicofollis dussumieri</i> (Valenciennes 1840)	Blacktip sea catfish	NE
	<i>Osteogeneiosus militaris</i> (Linnaeus 1758)	Soldier catfish	DD
	<i>Netuma thalassina</i> (Rüppell 1837)	Giant catfish	LC
Order: Syngnathiformes			
Mullidae	<i>Upeneus moluccensis</i> (Bleeker, 1855)	Goldband goatfish	LC
Order: Tetraodontiformes			
Balistidae	<i>Odonus niger</i> (Rüppell, 1836)	Red-toothed triggerfish	NE
	<i>Abalistes stellaris</i> (Bloch & Schneider 1801)	Starry triggerfish	LC
Tetraodontidae	<i>Takifugu oblongus</i> (Bloch, 1786)	Lattice blaasop	LC
Triacanthidae	<i>Triacanthus biaculeatus</i> (Bloch, 1786)	Short-nosed tripodfish	NE
	<i>Pseudotriacanthus strigilifer</i> (Cantor 1849)	Long-spined tripodfish	LC

Species of family Carangidae are of high commercial importance and also used as game fishes except *Atropus atropus*, *Alepes kleinii* and *Scomberoides tol* which are of less commercial importance. In the family Engraulidae, *Coilia mystus* forms the subsistence fisheries whereas others species are of commercial value. *Parascolopsis eriomma* of Nemipteridae also forms subsistence fisheries. Within Sciaenidae family, *Otolithoides biauritus* was of high commercial importance whereas *Otolithes ruber* was of minor commercial value; *Roncador stearnsii* was primarily used for game fishing. *Sardinella maderensis* of Clupeidae was highly exploited species in Gujarat. Out of all 114 species majority 71 were belonging to Least Concern (LC) category followed by Not evaluated (NE, $n = 16$), Data Deficient (DD, $n = 15$), Vulnerable (VU, $n = 4$), Near Threatened (NT, $n = 4$) and Endangered (EN, $n = 2$) categories (IUCN 2021). A total of 86 species were reported in a study on commercially important of marine finfish and shellfish of Okha, Dwarka district of Gujarat (Solanki *et al.* 2020). In another study (Joshi *et al.* 2018) 94 species were reported in Veraval of Gir-Somnath district. However, both studies have reported less number of species than the present study.

4 | CONCLUSIONS

The coast of Sutrapada supports rich ichthyofauna. This study confirmed presence of more species than other parts of Gujarat. Fishes are important, both commercially and ecologically, and therefore, the diversity of fishes needs to be conserved, valued and managed properly. The present study emphasizes on the proper management of the fishery and conservation aspects of the fishes at Sutrapada Coast.

ACKNOWLEDGEMENTS

The authors are grateful to the Head, Department of Zoology Faculty of Science, The Maharaja Sayajirao University of Baroda for the laboratory facilities and research work. One of the authors (SKS) is thankful to administration of Dalmia Public School, GHCL, Sutrapada. Authors would like to thank local fishermen for their valuable helps. Authors would also like to thank anonymous reviewers for their constructive comments.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

AUTHORS' CONTRIBUTION

SKS primary survey, collection of specimens and identification. **KJS** review of the manuscript and helped with the identification. **DMB** specimens collection, photography and manuscript preparation. **PCM** research supervision and review of the manuscript.

DATA AVAILABILITY STATEMENT

The data supporting the findings of this study are available within the article [and/or] its supplementary materials.

REFERENCES

- Balan K, Sivaraman P, George KP, Ramachandran M (1987) Appraisal of the marine fisheries of Gujarat. CMFRI Special Publication (38): 1–51.
- Day F (1878) The Fishes of India: being a natural history of fishes known to inhabit the seas and fresh waters of India, Burma and Ceylon. William Dawson and Sons Ltd., London. 778 pp.

- Day F (1889a) Fauna of British India, including Ceylon and Burma Vol. 2. Taylor and Francis, London 2: 1–509.
- Day F (1889b) Fauna of British India fishes. Taylor and Francis Ltd., London 1: 1–548.
- Eschmeyer WN, Fong JD (2014) Species by Family/Subfamily (<http://research.calacademy.org/research/ichthyology/catalog/SpeciesByFamily.asp>). Electronic version accessed on 18 May 2021.
- Frad, CMFRI (2018) Marine Fish Landings in India 2017. Technical report, CMFRI, Kochi. pp. 1–16.
- Fricke R, Eschmeyer WN, van der Laan R (Eds) (2020) Eschmeyer's catalog of fishes: genera, species, references. <http://researcharchive.calacademy.org/research/ichthyology/catalog/fishcatmain.asp>. Electronic version accessed on 18 May 2021.
- Galib SM (2015) Fish fauna of the Brahmaputra River, Bangladesh: richness, threats and conservation needs. *Journal of Fisheries* 3(3): 285–292.
- Galib SM, Naser SMA, Mohsin ABM, Chaki N, Fahad MFH (2013) Choice of fishes for consumption by the rural people of Bangladesh. *Trends in Fisheries Research* 2(1): 20–23.
- Galib SM, Rashid MA, Chaki N, Mohsin ABM, Joadder MAR (2016) Seasonal variation and community structure of fishes in the Mahananda River with special reference to conservation issues. *Journal of Fisheries* 4(1): 325–334.
- Galib SM, Samad MA, Mohsin ABM, Flowra FA, Alam MT (2009) Present status of fishes in the Chalan Beel—the largest Beel (wetland) of Bangladesh. *International Journal of Animal and Fisheries Science* 2(3): 214–218.
- Gopi KC, Mishra SS (2015) Diversity of marine fish of India. In: *Marine faunal diversity in India*, Academic Press. pp. 171–193.
- IUCN (2021) The IUCN red list of threatened species. Version 2021-1. <https://www.iucnredlist.org>, accessed on 18 May 2021.
- Jhingran VG (1983) Fish and fisheries of India, 1st edition. Hindustan Publishing Corporation, New Delhi. 660 pp.
- Joshi A, Parmar EAR, Temkar GS, Desai AY, Bhatt AJ (2018) [Ichthyofaunal biodiversity of Kharakuva fish market, Veraval, Gujarat, India](#). *International Journal of Bio-Resource & Stress Management* 9(5): 596–605.
- Mohanraj G, Nair KV, Asokan PK, Ghosh S (2009) Status of marine fisheries in Gujarat with strategies for sustainable and responsible fisheries. *Asian Fisheries Science* 22(1): 285–296.
- Mohsin ABM, Haque SMM, Galib SM, Fahad MFH, Chaki N, Islam MN, Rahman MM (2013) Seasonal abundance of fin fishes in the Padma River at Rajshahi district, Bangladesh. *World Journal of Fish and Marine Sciences* 5(6): 680–685.
- Mohsin ABM, Hasan MM, Galib SM (2009) Fish diversity of community based fisheries managed oxbow lake (Bookbhara Baor) in Jessore, Bangladesh. *Journal of Science Foundation* 7(1): 121–125.
- Murugan A, Namboothri N (2012) Finfishes of the Gulf of Mannar Biosphere Reserve: a field identification guide. Dakshin Foundation, Bangalore. 222 pp.
- Sarkar UK, Jena JK, Singh SP, Singh AK, Rebello SC (2012) Documenting coastal fish biodiversity of India: status, issues and challenges. In: *International Day for Biological Diversity Marine Biodiversity Uttar Pradesh State Biodiversity Board*. pp. 22–28.
- Sarwade JP, Khillare YK (2010) Fish diversity of Ujani wetland, Maharashtra, India. *The Bioscan* 1: 173–179.
- Solanki JB, Bajaniya VC, Parmar HV, Tank KV, Parmar HL (2020) Availability of commercially important of marine fin fish and shell fish along Okha fish landing centre, Gujarat. *Journal of Entomology and Zoology Studies* 8(1): 637–640.
- Talwar PK (1991) Inland fishes of India and adjacent countries. Vol. 1 and 2. CRC Press, Boca Raton.
- Zhang JB, Hanner R (2011) [DNA barcoding is a useful tool for the identification of marine fishes from Japan](#). *Biochemical Systematics and Ecology* 39(1): 31–42.



SK Singh  <https://orcid.org/0000-0002-4861-4242>

KJ Sarma  <http://orcid.org/0000-0002-5073-2539>

DM Bhatt  <https://orcid.org/0000-0003-2737-7951>

PC Mankodi  <https://orcid.org/0000-0002-8454-1984>